arXiv:gr-qc/0409001v3 13 Jan 2005

Euclidean action
for vacuum decay in a de Sitter universe

V. Balek* and M.Demetrian†

Department of Theoretical Physics, Comenius University, Mlynská dolina
842 48 Bratislava, Slovakia

February 7, 2008

Abstract

The behavior of the action of the instantons describing vacuum decay in a de Sitter is investigated. For a near-to-limit instanton (a Coleman-de Luccia instanton close to some Hawking-Moss instanton) we find approximate formulas for the Euclidean action by expanding the scalar field and the metric of the instanton in the powers of the scalar field amplitude. The order of the magnitude of the correction to the Hawking-Moss action depends on the order of the instanton (the number of crossings of the barrier by the scalar field): for instantons of odd and even orders the correction is of the fourth and third order in the scalar field amplitude, respectively. If a near-to-limit instanton of the first order exists in a potential with the curvature at the top of the barrier greater than 4 × (Hubble constant)^2, which is the case if the fourth derivative of the potential at the top of the barrier is greater than some negative limit value, the action of the instanton is less than the Hawking-Moss action and, consequently, the instanton determines the outcome of the vacuum decay if no other Coleman-de Luccia instanton is admitted by the potential. A numerical study shows that for the quartic potential the physical mode of the vacuum decay is given by the Coleman-de Luccia instanton of the first order also in the region of parameters in which the potential admits two instantons of the second order.

---

*e-mail address: balek@fmph.uniba.sk

†e-mail address: demetrian@sophia.dtp.fmph.uniba.sk

1 Introduction

The vacuum decay in a de Sitter universe has been considered as a mechanism of transition to a Friedman universe in the scenario of old inflation [1], and as a triggering mechanism in the scenario of open inflation [2]. In both scenarios, the instability of the false vacuum results in the formation of rapidly expanding bubbles containing the scalar field on the true-vacuum side of the potential barrier. The process is described by the Coleman-de Luccia (CdL) instanton [3]. Another kind of vacuum decay is mediated by the Hawking-Moss (HM) instanton [4] and proceeds in such a way that the field jumps on the top of the barrier inside a horizon-size domain. (For this interpretation, see [5].) Recently this process was proposed as a starting point for the inflation driven by the trace anomaly [6].

An important quantity characterizing the instanton is its action. It determines the probability of an instanton-induced transition per unit spacetime volume: if we denote the action by B, the probability is proportional to $\exp(-B)$. Thus, if more instantons exist for a given potential, the one with the least action prevails. In addition, the value of the action may be relevant to the question whether the instanton describes a well defined quantum mechanical transition at all. The transition takes place only if the instanton admits no negative modes of the perturbation superimposed on it; and according to [7], such modes are certainly not present only for the instanton with the least action. If negative modes exist, the preexponential factor in the expression for the probability diverges, which indicates that the wave function is not concentrated in a narrow tube around the instanton solution as it should be. (For an analogy in ordinary quantum mechanics, see [8].) Thus, the instanton with the least action, in addition to being the most probable, may be the only one with a physical meaning.

The values of the action of CdL instantons were discussed in the thin wall approximation in [3] and [9], and computed for a one-parametric class of quartic potentials admitting an instanton of the first order (an instanton with the scalar field crossing the top of the barrier just once) in [10]. Furthermore, the behavior of the action of CdL instantons was analysed in the approximation of negligible back reaction in [11]; the properties of a near-to-limit instanton (a CdL instanton close to some HM instanton) were studied with the back reaction included in [12]; and the instanton with the least action was identified among the instantons of odd orders in [7]. Here we link up to this research and develop it further. In sections 2 and 3 we find a formula for the action of a near-to-limit instanton with the back reaction included, and in section 4 we compute the instanton action for a class of quartic potentials with a subclass admitting instantons of the second order. Thus, in the first two sections we generalize the result of [11] and complete the results of [12], while in the last section we extend the results of [7] and [10].

2 The second order contribution to the action

Consider a compact O(4) symmetric 4D space and an O(4) symmetric scalar field ϕ living in it. The system is completely described by the functions $\phi(\rho)$ and $a(\rho)$, where ρ is the distance from the "north pole" (the radius of 3-spheres of homogeneity measured from the center) and a is the scale parameter (the radius of 3-spheres of homogeneity determined from the circumference). The compactness implies that ρ runs from zero to some ρ_f and $a = 0$ at both $\rho = 0$ and ρ_f. The Euclidean action is ($\hbar = c = 1$)

$$B = 2\pi^2 \int_0^{\rho_f} \left[\left(\frac{1}{2}\dot{\phi}^2 + V \right) a^2 - \frac{1}{\kappa}(\dot{a}^2 + 1) \right] a \, d\rho, \tag{1}$$

where the overdot denotes differentiation with respect to ρ, V is the effective potential of the scalar field and $\kappa = 8\pi G/3$. The extremization of the action yields

$$\ddot{\phi} + 3\frac{\dot{a}}{a}\dot{\phi} = \partial_\phi V, \quad \ddot{a} = -\kappa(\dot{\phi}^2 + V)a. \tag{2}$$

The equation for a is obtained by differentiating the equation

$$\dot{a}^2 = \kappa \left(\frac{1}{2}\dot{\phi}^2 - V \right) a^2 + 1,$$

and inserting for $\ddot{\phi}$ from the equation for ϕ. (In fact, when putting the variation of B equal to zero we obtain the first order equation for a with an additional term Ca^{-2}, where C is an integration constant. To suppress this term one has to introduce the lapse function into the expression for B and perform variation with respect to it.) Consider functions ϕ and a extremizing the action and such that the action is finite for them. Functions with these properties may be found, in principle, by solving equations (2) with the boundary conditions

$$\dot{\phi}(0) = \dot{\phi}(\rho_f) = 0, \quad a(0) = 0, \quad \dot{a}(0) = 1. \tag{3}$$

Since ϕ and $\dot{\phi}$ are finite at the points $\rho = 0$ and ρ_f, $\dot{a}$ and $\ddot{a}$ are finite at these points, too. Consequently, we may simplify the integral in (1) by replacing

$$a\dot{a}^2 \to -\frac{1}{3}a\dot{a}^2 - \frac{2}{3}a^2\ddot{a}$$

and using both equations for a to obtain

$$B = -\frac{4\pi^2}{3\kappa} \int_0^{\rho_f} a \, d\rho. \tag{4}$$

This expression is appropriate for numerical calculations but when expanding the action into the powers of the scalar field amplitude it is advisable to use expression (1) instead. Then one does not need to compute a up to the same order of magnitude as B; as we shall see, one gets along with the zeroth and second order contributions to a when computing the third and fourth order contributions to B, respectively.

Consider a potential with the global minimum, or the true vacuum, at $\phi = 0$, a maximum at $\phi = \phi_M > 0$ and a local minimum, or the false vacuum, at $\phi = \phi_m > \phi_M$. Suppose, furthermore, that the potential equals zero at the true vacuum. There are two trivial finite-action solutions to equations (2) with a compact 4-geometry,

$$\phi = \phi_M, \quad a = H_M^{-1}\sin(H_M\rho), \quad 0 < \rho < \pi H_M^{-1},$$

and

$$\phi = \phi_m, \quad a = H_m^{-1}\sin(H_m\rho), \quad 0 < \rho < \pi H_m^{-1},$$

where H_M and H_m are the values of $H = \sqrt{\kappa V}$ (the Hubble constant corresponding to a given value of ϕ) at $\phi = \phi_M$ and ϕ_m respectively. Both solutions consist of a 4-sphere and a constant scalar field. The first solution is the HM instanton; the second solution describes a "zero quantum transition" hence it may be regarded as a reference solution. In addition, the potential may admit nontrivial finite-action solutions to equations (2) called the CdL instantons. These solutions may be classified according to how many times the scalar field crosses the barrier: if the number of crossings is l, the instanton may be called "the CdL instanton of the lth order". Denote $\xi = \partial_\phi^2 V_M/H_M^2$. For $-\xi$ assuming one of the critical values

$$\lambda_l = l(l+3) = 4, 10, \dots \quad \text{for} \quad l = 1, 2, \dots, \tag{5}$$

the potential admits an approximate CdL instanton consisting of a 4-sphere with the radius H_M^{-1} and a scalar field proportional to the lth harmonic on the 4-sphere [13]. This solution may be called "the limit instanton of the lth order". If ξ is close to $-\lambda_l$, at least from one side, the potential should admit a CdL instanton close to the limit instanton with some small scalar field amplitude. In a one-parametric family of potentials, this near-to-limit instanton should approach the HM instanton as ξ approaches $-\lambda_l$.

Introduce dimensionless variables

$$\chi = H_M\rho, \quad u = \frac{1}{A}\Delta\phi, \quad v = H_M a,$$

where $\Delta\phi = \phi - \phi_M$ and $A = \Delta\phi(0)$, and write the potential as

$$V = H_M^2\left[\frac{1}{\kappa} + A^2\left(\frac{1}{2}\xi u^2 + \omega\right)\right].$$

The expression for the action, when rewritten into the dimensionless variables, is

$$B = \frac{2\pi^2}{H_M^2}\int_0^{\chi_f}\left[A^2\left(\frac{1}{2}u'^2 + \frac{1}{2}\xi u^2 + \omega\right)v^2 - \frac{1}{\kappa}(v'^2 - v^2 + 1)\right]v\, d\chi, \tag{6}$$

and the instanton equations are

$$u'' + 3\frac{v'}{v}u' - \xi u = \partial_u \omega, \quad v'' + v = -A^2\left(u'^2 + \frac{1}{2}\xi u^2 + \omega\right)v, \tag{7}$$

where $\chi_f = H_M \rho_f$ and the prime denotes differentiation with respect to χ. Denote $s = \sin\chi$ and $c = \cos\chi$. For the HM instanton $u = 0$, $v = s$ and $\chi_f = \pi$, which yields the action

$$B_M = -\frac{8\pi^2}{3\kappa H_M^2}. \tag{8}$$

For the limit instanton, u obeys the first equation (7) with $v = s$, $\xi = -\lambda_l$ and suppressed right hand side,

$$\mathcal{D}u = 0, \quad \mathcal{D} = \frac{d^2}{d^2\chi} - 3\text{cotan}\chi\,\frac{d}{d\chi} + \lambda_l.$$

The solutions are O(4) symmetric spherical harmonics in 5D or, up to normalization, the Gegenbauer polynomials with the parameter 3/2,

$$P_l = c,\ \frac{1}{4}(5c^2 - 1),\ \ldots \quad \text{for} \quad l = 1, 2, \ldots \tag{9}$$

When constructing a near-to-limit instanton one introduces expansions $\xi = \xi_0 + A\xi_1 + A^2\xi_2 + \ldots$, $u = Au_1 + A^2u_2 + A^3u_3 + \ldots$ and $v = v_0 + A^2v_2 + A^3v_3 + \ldots$ with $\xi_0 = -\lambda_l$, $u_1 = P_l$ and $v_0 = s$, and uses the expansion

$$\omega = \frac{1}{6}\eta A u^3 + \frac{1}{24}\zeta A^2 u^4 + \ldots,$$

where $\eta = \partial_\phi^3 V_M / H_M^2$ and $\zeta = \partial_\phi^4 V_M / H_M^2$. >From the expanded equations for u and v one obtains the functions u_1, u_2, ... and v_0, v_2, ..., and after inserting them into the expression for B one finds the coefficients B_0, B_2, ... in the expansion $B = B_0 + A^2B_2 + \ldots$ In particular, from $v_0 = s$ one obtains $B_0 = B_M$.

Let us prove that the term proportional to B_2 in the expansion of B vanishes so that $B = B_M$ up to the order A^2. For B_2 we have

$$B_2 = \frac{2\pi^2}{H_M^2}\int_0^\pi \left\{\frac{1}{2}(P_l'^2 - \lambda_l P_l^2)s^3 - \frac{1}{\kappa}[2scv_2' + (c^2 - 3s^2 + 1)v_2]\right\}d\chi. \tag{10}$$

(We have not included here a term arising from the shift of the upper limit of integration since such terms do not appear before the order A^4.) The first part, with the terms quadratic in P_l, may be rewritten using integration by parts to an integral of $P_l\mathcal{D}P_l$, hence it vanishes. The second part, with the terms linear in v_2, may be rewritten using integration by parts to an integral of $\mathcal{A}v_2$ with the coefficient of proportionality

$$\mathcal{A} = -(2sc)' + c^2 - 3s^2 + 1 = 0,$$

hence it vanishes, too. This can be seen also without an explicit calculation. The second part of B_2 is just what we obtain if we perform the variation of B_0 with respect to v_0 and put $\delta v_0 = v_2$, thus $\mathcal{A} = 0$ is in fact the equation determining v_0. As a result we obtain

$$B_2 = 0. \tag{11}$$

3 Higher order contributions to the action

The first higher order term in the expansion of B which may be nonzero is proportional to B_3. The terms in B_3 linear in u_2 and v_3 can be rewritten to be proportional to $\mathcal{D}P_l$ and $\mathcal{A}$, thus they vanish and we are left with the expression

$$B_3 = \frac{\pi^2}{H_M^2}\left(\xi_1 I_2 + \frac{1}{3}\eta I_3\right), \quad I_k = \int_0^\pi P_l^k s^3 d\chi. \tag{12}$$

To determine ξ_1, consider the equation for u_2

$$\mathcal{D}u_2 = \xi_1 P_l + \frac{1}{2}\eta P_l^2. \tag{13}$$

The task simplifies considerably by two observations: first, the left hand side, when expanded into the system of harmonics $P_0 = 1$, P_1, P_2, ..., does not contain a term proportional to P_l (this is easily seen if one expands u_2 into the harmonics and makes use of the fact that the operator $\mathcal{D}$, when applied on $P_{l'}$, reproduces it with a factor that is zero for $l' = l$); and second, the harmonics are orthogonal with the weight s^3. Thus, if we multiply equation (13) by P_l and integrate it with the weight s^3 we obtain zero on the left hand side. As a result we have

$$\xi_1 I_2 + \frac{1}{2}\eta I_3 = 0,$$

hence

$$B_3 = -\frac{\pi^2}{6H_M^2}\eta I_3.$$

The lth harmonic is a linear combination of c^l, c^{l-2}, ..., thus $P_l^3 s^3$ is an odd function on the interval $(0, \pi)$ and I_3 vanishes if l is odd. On the other hand, from the general formula for the integral of the product of three Gegenbauer polynomials [14] it follows that I_3 is nonzero if l is even. Thus, for even values of l both ξ_1 and B_3 are nonzero and the quantities $\Delta\xi = \xi + \lambda_l$ and $\Delta B = B - B_M$ are of the order A and A^3 respectively. Explicitly,

$$\Delta\xi \doteq -\frac{I_3}{2I_2}\eta A, \tag{14}$$

and

$$\Delta B \doteq -\frac{\pi^2 I_3}{6}\,\frac{\eta A^3}{H_M^2}. \tag{15}$$

The actual small parameter of the theory is not A but $\Delta\xi$, therefore we have to interpret the first equation as an approximate expression for A in terms of $\Delta\xi$. For $l = 2$, $I_2 = 2/21$ and $I_3 = 2/63$, thus the dimensionless coefficients in the expressions for $\Delta\xi$ and ΔB are $1/6$ and $\pi^2/189$ respectively. Note that we may avoid the computation of I_3 if we express B_3 in terms of ξ_1 and determine ξ_1 from the requirement that the term proportional to P_l on the right hand side of (13) vanishes.

To summarize, the correction to the HM action of order A^3 vanishes for instantons of odd orders, but it is nonzero for instantons of even orders provided η is nonzero. If we change the parameters of the potential so that ξ changes while η stays fixed, the character of the instanton changes as ξ crosses the value $-\lambda_l$. If, say, $\eta > 0$, for $\xi < -\lambda_l$ the function ϕ starts and ends to the right of ϕ_M (the instanton is "right handed"), while for $\xi > -\lambda_l$ the function ϕ starts and ends to the left of ϕ_M (the instanton is "left handed"). No matter what the handedness of the instanton, the action of the instanton is less than the HM action if $\xi < -\lambda_l$ and greater than the HM action if $\xi > -\lambda_l$.

To complete the analysis we have to compute B_4 for odd values of l. First, just as when we were computing B_3, we get rid of a large portion of the integrand by rewriting it in terms of $\mathcal{D}P_l$ and $\mathcal{A}$; in this way we remove the terms proportional to u_3 and v_4. Before we present the remaining terms let us mention a new point which arises in this order of expansion of B. One may expect that the value of v_2 at $\chi = \pi$ is nonzero and indeed, for $l = 1$ one finds $\delta \equiv v_2(\pi) = -\pi\kappa/8$ (see the appendix). Consequently, χ_f differs from π by a quantity of order A^2. As we shall see, this leads to additional terms in B_4 so that not the whole B_4 is stemming from the expansion of the integrand of B. After suppressing the terms proportional to u_3 and v_4 and integrating by parts we arrive at

$$B_4 = \frac{\pi^2}{H_M^2}\left\{\xi_2 I_2 + \frac{1}{12}\zeta I_4 + \int_0^\pi \left[u_2(-\mathcal{D}u_2 + P)s^3 + v_2\left(-\frac{2}{\kappa}\mathcal{B}v_2 + Q\right)\right] d\chi + b_4\right\}, \qquad (16)$$

where

$$P = \eta P_l^2, \quad Q = 3(P_l'^2 - \lambda_l P_l^2)s^2, \quad \mathcal{B} = -s\frac{d^2}{d^2\chi} - c\frac{d}{d\chi} - 2s,$$

and b_4 is a boundary term consisting of two parts, one coming from the shift in χ_f and the other coming from the integration by parts which has to be performed when deriving the term with the operator $\mathcal{B}$. The expression for B_4 simplifies further if we exploit the equations for u_2 and v_2,

$$\mathcal{D}u_2 = \frac{1}{2}P, \quad \mathcal{B}v_2 = \frac{1}{4}\kappa Q. \qquad (17)$$

Both equations are most easily derived by performing the variation of B_4 and using the fact that the operators $s^3\mathcal{D}$ and $\mathcal{B}$ are symmetric; however, they may be obtained by expanding the exact equations for u and v as well. The former equation is identical to (13) provided $\xi_1 = 0$, which is what we presently assume, and the latter equation is a linear combination of the two equations for v_2 presented in the appendix. If we make use of equations (17) and of the definition of P, we find that the integral in B_4 reduces to

$$\frac{1}{2}\eta J + \frac{1}{\kappa}\int_0^\pi Qv_2\ d\chi, \quad J = \int_0^\pi P_l^2 u_2 s^3\ d\chi.$$

Let us now show that the boundary term in B_4 is zero. We can find the correction to the upper limit of integration by noticing that the derivative of v_0 at $\chi = \pi$ is -1, so that to compensate for $\Delta v \doteq A^2 v_2$ in the neighborhood of $\chi = \pi$ we have to shift χ_f by $\Delta = A^2\delta$. The contribution to b_4 arising from this shift is

$$b_4^{shift} = A^{-4} \times \text{ the leading term in } \int_\pi^{\pi+\Delta} 2(F_0 + A^2 F_2)\, d\chi,$$

where F_0 and F_2 are the integrands in B_0 and B_2,

$$F_0 = \ldots - \frac{1}{\kappa}(c^2+1)s, \quad F_2 = \ldots - \frac{1}{\kappa}(c^2+1)v_2.$$

(The omitted terms are irrelevant for the present discussion.) The leading term in the integral of F_0 equals $F_0'(\pi)\Delta^2/2 = \Delta^2/\kappa$, since $F_0(\pi) = 0$, and the leading term in the integral of F_2 equals $F_2(\pi)\Delta = -2\delta\Delta/\kappa$, hence

$$b_4^{shift} = -\frac{2\delta^2}{\kappa}.$$

On the other hand, from the expression of the relevant part of B given in the appendix it follows that the contribution to b_4 coming from the integration by parts is

$$b_4^{int} = \frac{2\delta^2}{\kappa}.$$

Putting this together we find

$$b_4 = 0. \tag{18}$$

Next we have to write down the equation for u_3 in order to determine ξ_2. When constructing this equation we encounter a subtlety that is again connected with the presence of δ. If δ is nonzero, v_2 is not small with respect to s in the vicinity of $\chi = \pi$ and the factor $\dot{a}/a = H_M v'/v$ in the equation for ϕ cannot be treated perturbatively. In fact, if we formally expand v'/v up to the order A^2,

$$\frac{v'}{v} \doteq \text{cotan}\chi + A^2 q, \quad q = \left(\frac{v_2}{s}\right)',$$

we can see that the correction term even diverges relatively to the zeroth order term in $\chi = \pi$. (It behaves like $-\Delta/\epsilon^2$ for $\epsilon = \pi - \chi \to 0$, while the zeroth order term behaves like $1/\epsilon$.) To fix that, note that the term $\hat{\delta}$ in v_2 which is nonzero at $\chi = \pi$ and therefore is responsible for the nonzero value of δ equals $-\delta\chi v_0'/\pi$ (see the appendix), thus $\hat{\delta}$ may be absorbed into v_0 by redefining $\chi \to (1 - \Delta/\pi)\chi$. Such procedure is well known in the perturbation theory of an anharmonic oscillator where the terms of the type $\hat{\delta}$ are called "resonance terms" [15]. The corrected equation for v_3 is obtained in such a way that we replace q by q^{red} ("red" standing for "reduced"), defined as q with v_2 replaced by $v_2^{red} = v_2 - \hat{\delta}$, and add to ξ_2 the term κ arising from the redefinition of χ in the equation for u_1. However, these corrections, necessary as they are when one computes v_3, are superfluous if one is interested only in the integral of the equation for v_3 with the weight s^3. The point is that the weight washes out the singularity in q, therefore the reverse transformation $\chi \to (1 + \delta/\pi)\chi$ by which one passes from the corrected equation to the original one may be treated on equal footing with the regular transformations (that is, transformations that do not move the point $\chi = \pi$). To confirm this, one may check that the two corrections mentioned above cancel when the equation is integrated over χ with the weight s^3. The uncorrected equation for v_3 reads

$$\mathcal{D}v_3 = \xi_2 P_l + \eta_1 P_l u_2 + \frac{1}{6}\zeta P_l^3 - 3qP_l', \tag{19}$$

and by the same argument as we have used when calculating B_3 we can deduce from it the identity

$$\xi_2 I_2 + \eta J + \frac{1}{6}\zeta I_4 + \int_0^\pi 3qP_l P_l' s^3 \, d\chi = 0.$$

Finally, by integrating by parts and using the equation for P_l it may be shown that the integrand in the last term can be replaced by $-Qv_2$. This completes the proof that in our initial expression for B_4 the sum of all terms following $\xi_2 I_2$ equals $-\xi_2 I_2/2$, and

$$B_4 = \frac{\pi^2}{2H_M^2} I_2 \xi_2.$$

Similarly as when we were computing B_3 we can now obtain approximate expressions for $\Delta\xi$ and ΔB in terms of A. Again, we may determine ξ_2 also by analysing the equation for v_3; however, for that purpose the corrected equation has to be used, with q replaced by q^{red} and ξ_2 by $\xi_2 + \kappa$. Let us present the results for the most important case $l = 1$. In this case,

$$\Delta\xi \doteq -\frac{1}{14}\left(\frac{1}{12}\eta^2 + \zeta + 32\kappa\right) A^2 \tag{20}$$

and

$$\Delta B \doteq \frac{2\pi^2}{15} \frac{\Delta\xi A^2}{H_M^2}. \tag{21}$$

The expression for $\Delta\xi$ implies that a near-to-limit instanton exists only if $-\xi$ approaches the value $\lambda_1 = 4$ from a given side. Which side, that depends on the parameters η and ζ. Define

$$\zeta_{crit} = -\frac{1}{12}\eta^2 - 32\kappa. \tag{22}$$

If $\zeta > \zeta_{crit}$ the instanton exists for $\xi < -4$, while if $\zeta < \zeta_{crit}$ the instanton exists for $\xi > -4$. The action of the instanton is less than the HM action in the former case and greater than the HM action in the latter case.

The expression for $\Delta\xi$ is identical to that obtained in [12], and if we suppress κ in the expression for $\Delta\xi$ and insert for H_M, η and ζ the values for the quartic potential, the resulting expression for ΔB is identical to that derived in [11]. Our formula for ΔB is consistent also with the behavior of the first perturbation mode of the near-to-limit instanton established in [12], since this mode is positive (and, consequently, contributes to the action by a positive quantity) in the same range of parameters in which we found that $B_M - B$ is positive.

4 The instanton with the least action

The question which instanton has the least action is addressed in [7] where it is shown that among the instantons of odd orders, the one with the least action is necessarily of the first order. The proof is based on the analysis of a graph that may be regarded as a phase diagram for the solutions of the Euclidean theory. One introduces *noninstanton* solutions $\phi_-(\rho)$ and $\phi_+(\rho)$ starting to the left and to the right of ϕ_M, and assigns to them the values of $\Delta\phi$ and $\pi_\phi = 2\pi^2 a^3 \dot{\phi}$ at such ρ at which the function $a(\rho)$ reaches maximum. In this way one obtains two oriented curves in the $(\Delta\phi, \pi_\phi)$ plane: the curve C_- starting at the point $(-\phi_M, 0)$ (the true vacuum) and ending at the origin (the top of the barrier), and the curve C_+ starting at the origin and ending at the point $(\phi_m - \phi_M, 0)$ (the false vacuum). An instanton of an odd order is represented by an intersection of the curves C_+ and C_-, and its "net action" ΔB equals the area inside a loop passing from the origin to the intersection and back, first along the curve C_+ and then along the curve C_-. (The area is regarded here as the z-component of the vector $\oint \mathbf{r} \times d\mathbf{r}$, where $\mathbf{r}$ is the radius vector in the (x, y) plane. According to this definition, the area inside a given curve is positive if the curve is oriented counterclockwise and negative if the curve is oriented clockwise.) As to the instantons of even orders, they are represented by the points where the curves C_- (for left handed instantons) and C_+ (for right handed instantons) intersect the axis $\Delta\phi$, and their net action equals twice the area inside a loop passing from the origin to the intersection and back, first along the curve C_- and then along the axis $\Delta\phi$ if the instanton is left handed, and first along the axis $\Delta\phi$ and then along the curve C_+ if the instanton is right handed. Three typical phase diagrams are depicted in fig. 1. All diagrams contain an instanton of the first order denoted by 1;



Figure 1: Phase diagrams for the solutions of the Euclidean theory

besides, the middle diagram contains two left handed instantons of the second order denoted by $2'_-$ and $2''_-$, and the right diagram contains one left handed and one right handed instanton of the second order denoted by 2_- and 2_+. The net action of the instanton of the first order equals the area of the light grey patch in the left diagram; the area of the the light grey patch and the lower dark grey patch in the middle diagram; and the area of the light grey patch and both dark grey patches in the right diagram. The limit curves are in all three cases oriented clockwise, therefore the area is negative and the action of the instanton is less than the HM action. The net action of the instantons of the second order equals twice the area of the corresponding dark grey patches; consequently, the net action is negative for both 2_- and 2_+, positive for $2'_-$, and either negative or positive, depending on which of the two dark grey patches prevails, for $2''_-$. There seems to be no general argument restricting the area of the dark grey patches in comparison with the light grey one, therefore the less of the actions of the two instantons of the second order may be also less than the action of the instanton of the first order. As seen from the geometry of the patches, one can only state that the action of the instanton of the first order is less than the *mean* action of the instantons of the second order.

The phase diagrams suggest that if we take into consideration the instantons of even orders (including the HM instanton which may be regarded as the instanton of the zeroth order), the instanton with the least action may not stay of the first order. We have seen that if two instantons of the second order exist in addition to one instanton of the first order, the instanton with the least action may one of them. In a similar way one may prove that if there are two instantons of the first order and none of a higher order (which occurs if the curves C_- and C_+ are intertwined next to the origin and do not intersect the axis $\Delta\phi$ somewhere), and if both instantons have approximately the same action (which occurs if the two intersections of the curves C_- and C_+ outside the origin are close to each other), the instanton with the least action is necessarily the HM instanton. For a typical potential, however, one expects the instanton of the first order to prevail even in the presence of instantons of even orders. To support this opinion, we present here a numerical study for a certain class of quartic potentials.

The quartic potential is usually regarded as depending on three parameters, however the parameter m whose square appears in front of ϕ^2 can be removed from the theory if one makes use of rescalings $a \to a/m$ and $\rho \to \rho/m$ in the equations for ϕ and a. The rescaled potential is

$$V = \frac{1}{2}\phi^2 - \frac{1}{3}\delta\phi^3 + \frac{1}{4}\lambda\phi^4. \tag{23}$$

The potential has the desired form only for the parameter δ running from $\delta_m = 2\sqrt{\lambda}$ to $\delta_M = 3\sqrt{\lambda/2} \doteq 1.06\delta_m$. For numerical calculations we have chosen the value of δ in the middle of this interval,

$$\Delta \equiv \frac{\delta - \delta_m}{\delta_M - \delta_m} = 0.5.$$

The parameter ξ may be expressed in terms of δ and λ as $(G = 1)$

$$\xi = -\frac{9\lambda}{\pi}\,\frac{X(1+X)}{(1-X)(1+3X)}, \quad X = \sqrt{1-\frac{4\lambda}{\delta^2}}. \tag{24}$$

Note that X depends on δ and λ only through Δ, so that if one keeps Δ fixed ξ becomes proportional to λ.

The results are summarized in fig. 2 and 3. The characteristics of instantons obtained numerically by solving the equations (2) are depicted by solid lines, and those computed from the formulas (20) and (21) in the neighborhood of $\xi = -4$, and from the formulas (14) and (15) in the neighborhood of $\xi = -10$, are depicted by dotted lines. As ξ decreases, the phase diagram passes through the three regimes visualised in fig. 1. They occur, from the left to the right, for ξ running from -4 to some ξ_{crit} greater than -10; for ξ running from ξ_{crit} to -10; and for ξ running from -10 to -18. The behavior of instantons near the limit values of ξ is determined by the fact that for the quartic potential both η and ζ are positive. As a result, the first order instantons exist only for $\xi < -4$ and the near-to-limit second order instantons are right handed for $\xi < -10$ and left handed for $\xi > -10$.

In fig. 2 the limit values of ϕ are plotted as functions of the parameter ξ. The graph to the right provides a magnified view of a narrow segment of the graph to the left. For comparison, the location of the top of the barrier and of the false vacuum is shown in the graph to the left.



Figure 2: Range of values of the scalar field for instantons in the quartic potential

In fig. 3 the ratio $\Delta B/B_M$ is plotted as a function of the parameter ξ. Again, we can see a tiny segment of the graph to the left displayed in more detail in the graph to the right. The



Figure 3: Action of instantons in the quartic potential

values of $\Delta B/B_M$ in the latter graph are to be multiplied by 10^{-4}. The HM action is negative, hence in the region above the horizontal axis there holds $\Delta B < 0$.

5 Conclusion

The behavior of the action of CdL instantons has been investigated in two complementary ways, analytically for a general potential with the value of ξ (the rescaled second derivative of V at $\phi = \phi_M$) near to one of the limit values $-4, -10, -18, \ldots$, and numerically for a class of quartic potentials with the value of ξ running from -4 to -18. The analytical study implies that the properties of instantons of odd and even orders are in many respects different. While the instantons of even orders exist for ξ on both sides of the limit value, the instantons of odd orders exist only for ξ either to the left or to the right of the limit value, depending on the values of η and ζ (the rescaled third and fourth derivative of V at $\phi = \phi_M$). Also the dependence of the action on the scalar field amplitude is different for instantons of even and odd orders. However, the two kinds of instantons have one property in common: their action is less than the HM action if the value of ξ is less than the limit value, that is if the potential is more curved at the top of the barrier than the potential for which the near-to-limit instanton shrinks to the top of the barrier, and vice versa. (Note that this may be proven also without an explicit computation, by investigating the properties of the phase diagram in the neighborhood of the origin.) As may be shown by the analysis of noninstanton solutions [13], if a potential which admits a near-to-limit instanton is less curved than the limit potential, it necessarily admits another instanton of the same order, with the scalar field extended along a finite segment of the barrier. The analysis of the phase diagram as well as numerical calculations performed for the second order instantons suggest that the extended instanton has less action than the compact one, but as both instantons first appear in the course of decreasing of ξ they have greater action than the HM instanton. The crucial question is which instanton has the least action. We have shown that (1) for potentials with $\xi \lesssim -4$ and $\zeta < \zeta_{crit}$, the instanton with the least action is the CdL instanton of the first order if the potential admits no other CdL instanton, (2) for quartic potential with $\Delta = 0.5$ and $\xi < -4$, the same is true also if the potential admits two instantons of the second order. Note that the order of the instanton determines the number of bubbles and walls filled with the scalar field on the true-vacuum side of the barrier which are created in the process of quantum mechanical tunneling. For an instanton of the first order the outcome of the tuneling is one bubble and no wall, while for instantons of the second order the outcome is either no bubble and one wall or two bubbles and no wall. Thus, in both cases cited above the vacuum decays via formation of a single bubble.

Acknowledgement. This work was supported by the grant VEGA 1/0250/03.

A The second order contributions to u and v

To derive the expression (20) for $\Delta\xi$ we need to know the functions u_2 and v_2. The equation for u_2 solves for $l = 1$ by the Ansatz $u_2 = C_1 + C_2 c^2$. (Our definition of A requires that $u_2(0) = 0$, which can be fulfilled only if we include into u_2 also a term proportional to c. However, since this term may be absorbed into u_1 it has no effect on the results and may be ignored.) The solution is

$$u_2 = \frac{A^2}{12}\eta\left(\frac{1}{2} - c^2\right). \tag{A-1}$$

The equation for v_2 obtained from the second order equation for v reads

$$\mathcal{B}^{(2)} v_2 = -\kappa(P_l'^2 - 2P_l^2)s, \quad \mathcal{B}^{(2)} = \frac{d^2}{d\chi^2} + 1. \tag{A-2}$$

This is supplemented by the initial conditions $v_2 = v_2' = 0$ at $\chi = 0$. For $l = 1$, the right hand side is proportional to $(s^2 - 2c^2)s = (1 - 3c^2)s$, which suggests the Ansatz $v_2 = (C_1 + C_2 c^2)s + C_3 \chi c$. (The term proportional to c is omitted because of the first initial condition.) If we insert this Ansatz into the equation for v_2 and use the second initial condition we obtain

$$v_2 = \frac{1}{4}\left[\left(1 - \frac{3}{2}c^2\right)s + \frac{1}{2}\chi c\right]. \tag{A-3}$$

This provides the expressions for δ and $\hat{\delta}$ cited in section 3. The latter expression is valid also for higher values of l since v_2 always splits into a part containing s and a term proportional to χc. Note that our expressions for u_2 and v_2 are consistent with the expansions of a and ϕ in [12].

The expression (16) for B_4 allows us to avoid an explicit computation of this quantity once we have computed $\Delta\xi$. For completeness, let us explain how the part of (16) containing v_2 is obtained. The starting expression is

$$\int_0^\pi \left[Qv_2 - \frac{2}{\kappa}(2cv_2v_2' + s{v_2'}^2 - 3sv_2^2)\right] d\chi.$$

Rewrite the expression in the round brackets as

$$(cv_2^2)' + (sv_2v_2')' - 2sv_2^2 - (sv_2')'v_2.$$

The integral of the first two terms over χ equals $-\delta^2$ which leads to the expression for b_4^{int} cited in section 3, while the remaining two terms may be rewritten as $v_2\mathcal{B}v_2$. The equation for v_2 resulting from the variation of B_4 (the second equation (17)) can be obtained also by combining the first order equation for v_2,

$$\mathcal{B}^{(1)}v_2 = -\frac{1}{2}\kappa\left(\frac{1}{2}{P_l'}^2 + 2P_l^2\right)s^2, \quad \mathcal{B}^{(1)} = c\frac{d}{d\chi} + s, \qquad \text{(A-4)}$$

with the equation (A-2). For that purpose, note that $\mathcal{B} = -s\mathcal{B}^{(2)} - \mathcal{B}^{(1)}$.

References

[1] A. H. Guth, Phys. Rev. **D23**, 347 (1981).

[2] M. Bucher, A. S. Goldhaber and N. Turok, Phys. Rev. **D52**, 3314 (1995); K. Yamamoto, M. Sasaki and T. Tanaka, Astrophys. J. **455**, 412 (1995); A. D. Linde, Phys. Lett. **B351**, 99 (1995); A. D. Linde and A. Mezhlumian, Phys. Rev. **D52**, 6789 (1995).

[3] S. Coleman and F. de Luccia, Phys. Rev. **D21**, 3305 (1980).

[4] S. W. Hawking and I. G. Moss, Phys. Lett. **110B**, 35 (1982).

[5] A. D. Linde: *Particle Physics and Inflationary Cosmology*, Harwood, Chur, Switzerland (1990).

[6] S. W. Hawking and T. Hertog, Phys. Rev. **D66**, 123509 (2002).

[7] T. Tanaka, Nucl. Phys. **B 556**, 373 (1999).

[8] T. Banks, C. Bender and T. T. Wu, Phys. Rev. **D8**, 3346 (1973); 3366 (1973).

[9] S. Parke, Phys. Lett. **B121**, 313 (1983).

[10] D. A. Samuel and W. A. Hiscock, Phys. Rev. **D44**, 3052 (1991).

[11] M. Sasaki, E. D. Stewart and T. Tanaka, Phys. Rev. **D50**, 941 (1994).

[12] T. Tanaka and M. Sasaki, Progr. Theor. Phys. **88**, 503 (1992).

[13] L. Jensen and P. J. Steinhardt, Nucl. Phys. **B237**, 176 (1984); V. Balek and M. Demetrian, Phys. Rev. **D69**, 063518 (2004).

[14] N. J. Vilenkin, *Special Functions and the Theory of Group Representations*, Providence, RI: Amer. Math. Soc. (1968).

[15] L. D. Landau and E. M. Lifshitz, *Mechanics (Course of Theoretical Physics)*, Butterworth-Heineman (1995).